

ORGANIC COMPOSITIONS AND METHODS OF USE FOR PROMOTING PLANT GROWTH

FIELD OF THE INVENTION

5 The invention relates to new compositions and methods of increasing valuable traits in agronomically important plants. Compositions of the present invention are formulated from natural organic components including seaweed and fish hydrolysates. Cold-tolerant seaweed species, for example, are preferred for use in compositions of the present invention. Geothermal water, high in sulfur and minerals, is preferred for use in preparing compositions
10 of the present invention by means of bio-processing fermentation. Formulations of the present invention can be applied effectively to a wide variety of plant species to improve valuable agronomic traits including but not limited to growth and yield. The stimulation of the production of phenolic metabolites and aromatic compounds, for example, in treated plants leads to a significantly perceived improvement in the flavor and natural aroma of
15 different crops. A relatively small amount of the bioprocessed compositions described herein are required to be effective in enhancing valuable traits in agronomic crops, thus an affordable advantage is provided, particularly to organic growers.

BACKGROUND OF THE INVENTION

20 The use of botanical extracts such as seaweed in organic fertilizers has been well known for years and has been successfully developed in the USA and Europe. It is also known that proline is beneficial for improving shoot organogenesis. A natural source of proline and proline precursors can be obtained from fish protein hydrolysates (FPH), a byproduct of the fishery industry. FPH is employed in compositions described herein to stimulate proline
25 synthesis and shoot organogenesis by exploiting the proline-linked pentose phosphate pathway (PPP). In the presence of elevated levels of proline, stimulation of cytokinins and auxin occurs via the PPP and shikimate pathways, respectively. Treatments with FPH significantly increases the endogenous proline content and the extent of development. See, *e.g.*, Milazzo MC, Shetty K., *et al.*, J. Agric. Food Chem., 47(4):1771 (1999).

30

Genetically modified plants and produce engineered to enhance value-added traits are generally not well-accepted by buyers and consumers in industrialized nations. Moreover, consumers are generally more receptive to organically grown fruits, vegetables and plants in

general. Accordingly, a cost-effective solution to meeting the demand for organically grown produce having enhanced agronomic traits would be valuable indeed.

SUMMARY OF THE INVENTION

5 The present invention is directed to organic compositions and methods for increasing valuable agronomic traits in plants. Particularly, organic compositions of the present invention are comprised of an aqueous solution of seaweed extract and fish hydrolysates. Preferred compositions are comprised of an extract of at least one cold-tolerant seaweed species. Preferred embodiments of the present invention are formulated with geothermal
10 water which is high in sulfur and minerals. In addition, the current invention is directed to a method of preparing organic compositions for increasing valuable agronomic traits in plants comprising mixing seaweed extract with fish hydrolysates and fermenting the mixture with geothermal water. The invention is further directed to methods of applying organic compositions of the present invention to increase valuable agronomic traits in plants.

15

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 displays improved vigor of pea seedlings. Left: control. Right: treated at a dose of 2 ml organic composition of the present invention per liter of water.

20 **Figure 2** shows improved vigor of Rosemary plants. Left: Rosemary plants soil treated with a drench of organic composition of the present invention at a dose of 5 ml/liter. Right: control

Figure 3 illustrates enhanced vigor of corn seeds treated with 2 ml organic composition of the present invention /liter of water (right). Left is the control.

25 **Figure 4** illustrates Shoot and Root enhancement in tomato seedling on post-germination day 12. A. Control; B. Following treatment at a dose of 2.5 ml organic composition of the present invention /liter water; C. Following treatment at a dose of 5 ml organic composition of the present invention /liter water.

Figure 5 displays a graph illustration of the results of FIG.4.

DETAILED DESCRIPTION OF THE INVENTION

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. All publications and patents referred to herein are incorporated by reference.

5

Compositions for increasing valuable agronomic traits in plants are provided comprised of seaweed extract and fish hydrolysate. Methods of preparing compositions of the invention are provided which comprise mixing seaweed extract with fish hydrolysate and fermenting the mixture with Icelandic geothermal water. Methods of increasing at least one valuable
10 agronomic trait in a plant are provided comprising applying a composition comprising seaweed extract and fish hydrolysate to the plant. U.S. Patent No. 5,906,941, entitled, *Plant Propagation Compositions and Methods* is incorporated herein by reference.

Organic compositions of the present invention are derived from natural sources. Particularly
15 described herein are bioprocessed compositions originating from natural marine and botanical extracts for increasing valuable agronomic traits in plants. Compositions of the present invention are a formulated aqueous solution of seaweed extract and fish hydrolysates. Compositions described herein are high in soluble amino acids to induce the proline pathway which is a critical developmental mechanism in plants directly related to valuable agronomic
20 traits. Organic compositions of the present invention stimulate proline synthesis, shoot and root vigor and shoot organogenesis by exploiting the proline-linked pentose phosphate pathway (PPP). The presence of elevated levels of proline confers stimulation of cytokinins and auxin occurs via the PPP and shikimate pathways, respectively. Treatments with compositions of the present invention significantly increase the endogenous proline content
25 and the extent of plant root and shoot development and strength. Compositions of the present invention also contain natural plant growth promoting substances including, for example, cytokinins, auxins and amino acids such as proline, glutamic acid and arginine. Formulations described herein are used effectively on a wide variety of plant species to improve growth and yield. A relatively small amount of the bioprocessed composition is required to enhance
30 valuable traits in agronomic crops. Accordingly, an affordable advantage is provided particularly to organic growers.

Fish and seaweed for use in preparing organic compositions (fermented product) of the present invention are preferably harvested from cold waters of the North Atlantic, e.g., off the coast of Iceland. However, compositions of the present invention may be prepared using a myriad of ingredients similar to those described herein, e.g., fish, seaweed, and water from
5 around the world.

The world annual fish catch has been estimated to be approximately 71 million tons, only about 25 tons of which is used for human food. About 20 million tons is caught specifically to be made into fish meal. Of the remaining 26 million tons of "waste", about half is recovered
10 for fish meal, while the rest is made up of bones and offal. Mackie, Process Biochem., 17:26-31 (1982). This solid waste is high in protein. Fishery by-products, i.e., soluble fish protein hydrolysate (SFPH) are made from the solid waste left behind after fish processing. See, U.S. Patent No. 5,906,941, entitled *Plant Propagation Compositions and Methods*, incorporated herein by reference.

15

Fermenting fish to produce a soluble fish protein hydrolysate (SFPH)

Compositions of the present invention are generally comprised of slightly acidified amino acids such as glutamic acid, arginine and proline obtained from fermentation of fish to prepare hydrolysates. Fish hydrolysates are particularly preferred from capelin (and other
20 pelagic fish species), herring and menhaden and are enriched by means of the process described herein in glutamic acid, arginine, and proline. The first stage of the process of preparation of compositions of the present invention is the initial stage of fermentation wherein fish and/or portions or remains after processing ("fish") are added to a tank (grinding or physical mincing of the fish is optional). For example, 1000kg of fish may be added to a
25 1200 liter tank at room temperature (~20-26°C), e.g., 23°C. Temperature, however, is not a critical aspect of the process of the invention. Indeed, processes described herein may be carried out well below or well above room temperature. The temperature at which the process is carried out merely affects an amount of time otherwise required for fermentation. The initial stage of fermentation is preferably, although not necessarily, enhanced by the
30 addition of about, for example, 3% to about 5% v/w lactic acid to the tank holding the fish. For example, to a tank holding about 1000kg fish about 25 to about 60ml lactic acid may be added to enhance the fermentation process. Lactic acid is a by-product of the dairy industry and is readily available from many different commercial sources. Alternative acids to lactic

acid that may similarly be employed to enhance this fermentation process include, for example, citric acid, acetic acid, and malic acid. However, *in addition* to lactic acid (or citric acid, acetic acid, or malic acid) the addition of about 1% to about 2% v/w formic acid is also preferably employed to enhance the fermentation process. Industrial grade formic acid is
5 available from a wide variety of commercial sources. The initial fermentation process is generally allowed to proceed in the tank for about three (3) to about ten (10) days before the addition of seaweed extract. About four (4) to five (5) days, for example, is a usual initial incubation period of fish fermentation at room temperature for the production of SFPH.

10 Homogenized Seaweed

Seaweed extracts, for example, from cold-tolerant Icelandic species are preferred for use in the preparation of compositions of the present invention. Extracts from cold-tolerant seaweed species, e.g., the genus *Laminaria* (PHEOPHYCEES) (e.g., *Laminaria hyperborea*, *Laminaria digitata* and *Laminaria nigripes*) may be used. Seaweed extract from *Laminaria* species
15 growing in cold water generally contain proline in the range of 0.1-0.5% and phenolics in the range of 0.1%-0.25%. These same extracts produce 0.01% to 0.05% cytokinins. Accordingly, seaweed extracts employed in compositions of the present invention generally provide a source of cytokinins which contribute significantly to the stimulation of shoot growth. Cold-tolerant (e.g., Icelandic) species also, for example, have natural stress
20 modulators from the proline pathway.

Seaweed is collected and preferably dried. Heat dried seaweed is preferred which has about 10-20% water content. The texture is approximately that of paper. The seaweed is powdered. Although particle size is not intended to be a limitation of the claims appended
25 hereto, an average particle size of crushed and/or “blended” seaweed may be between about 0.5mm to about 5mm. Seaweed may be, for example, homogenized into a paste using a blender. The term homogenized, as used herein, however, refers to all of the above descriptions in this paragraph. The seaweed extract for use in the method described herein is preferably prepared by combining about 10% to about 20% dried/ground seaweed with water
30 w/v. Geothermal water relatively high in sulfur, for example, and minerals (e.g., Icelandic), is preferred.

Bio-processing fermentation

Seaweed extract described herein is then added to the initial fermentation (SFPH production) process described *supra*. Homogenized seaweed is added to the SFPH to produce a volume so that at least about 1% but less than about 20% of the volume is initially due to seaweed.

- 5 The term seaweed, as used herein, generally refers to seaweed, *per se*, including all cell contents. Generally, about 10% to about 25% seaweed extract v/v is added to the fermenting SFPH. Accordingly, for example, to accomplish 20% v/v seaweed extract, about 200 liters seaweed extract is added to about 800 liters initially fermenting SFPH. The volume is then fermented to hydrolyse the seaweed and to further hydrolyse the SFPH. Bio-processing
- 10 (fermentation) is continued for about five (5) to about twenty (20) *more* days (past step 1, the initial stage of fermentation or initial SFPH preparation). About ten (10) days is preferred at room temperature (~20-26°C), e.g., 23°C. However, temperature is not a limitation. Organic acid and natural fermenting extracts of fish are allowed to ferment and naturally disassociate and hydrolyse the fish and seaweed in natural high sulfur geothermal water. When the period
- 15 of bio-processing fermentation of the SFPH/seaweed extract (preferably prepared with geothermal water) is substantially complete, the fermenting mixture is *preferably* - stirred and transferred to separate container(s) (e.g., drums) - and allowed to gravity-settle for about five (5) to about ten (10) days (alternatively, to speed the separation aspect of the process, the mixture could be subject to centrifugation). The bio-processing fermentation mixture is
- 20 generally separated into three (3) layers or phases. The resulting top layer (soluble upper phase), about 40% of the volume, is the primary organic aqueous liquid composition of the present invention (referred to herein as the organic composition of the present invention). The resulting top layer, which is the organic composition for increasing valuable agronomic traits in plants, is separated from the volume. The "organic composition" is thereby separated
- 25 and bottled. The lower solid phase may be blended with other solid organic compost material, for example, to make solid fertilizer.

- Among the important amino acids enriched in the resulting organic composition of the present invention are the acidified amino acids, e.g., glutamic acid (generally 6% -10% of
- 30 soluble nitrogen), arginine (3-5% of soluble nitrogen), and proline (2-4% of soluble nitrogen). In addition, phenolics result in the range of 0.1%-0.2%. The sulfur content, from geothermal water, is also enriched from, for example, 0.05%- 0.1%. Acidified amino acids are key organic stimulators of the stress modulating proline-linked pentose phosphate

pathway. Acidified amino acids provide benefits, for example, to stimulate soil conditioning micro-organisms and plant promoting micro-organisms. Fish extracts employed in compositions of the present invention provide small fatty acids that are anti-microbial against acid intolerant plant bacterial pathogens.

5

Bio-processing fermentation, for example, using geothermal water (e.g., Icelandic) high in sulfur and minerals and the enhanced stress modulating natural compounds derived from seaweed together with the fish extracts described herein, provides key advantages.

Accordingly, preferred water for use in fermenting compositions of the present invention is

10 geothermal water enriched in sulfur, e.g., from Icelandic geothermal sources.

Liquefaction is carried out by enzymes already present in the fish. Liquefaction is entirely due to autolysis which can be enhanced by temperature control. At higher temperature of 20-30°C liquefaction occurs rapidly. However, this process is slower at lower temperatures.

15 See, e.g., Tatterson, I.N., *et al.*, J. Sci. Food Agric., 25: 369-379 (1974). Acids may be added to provide favorable conditions for enzyme activity and inhibition of spoilage bacteria.

Formic acid of about 3% has been used in many studies with good results. However, citric acid and lactic acid from beverage and milk processing can be used effectively from 0.5% to 5%.

20

The acidic treatment (lactic and formic acid) is performed in a blend with Icelandic seaweed with Icelandic geothermal water high in sulfur. This is uniquely targeted to caplein fish species at low levels of fat content (1%-5%) after spawning and, for example, can be extended to other pelagic fish species such as menhaden and herring. Once the fish silage-seaweed is made from sulfur geothermal water, it becomes enriched in amino acids glutamic acid, proline and arginine. Caplein species, for example, about 1000kg or its multiples are hauled by fishing boats from coastal Iceland and added to tanks ranging from 1000kg capacity to 12,000 kg capacity. The fish can be ground in some cases before adding. The moisture content of fish is about 75%-80% and ash content of about 3.5-4.5%. The oil content at this stage ranges from 1%-5% (low fat) following spawning. The total nitrogen content is about 1.7% to 2.0 %. Fresh fish is added to tanks with and without grinding and a combination, for example, of 2%-3% lactic acid from dairy processing waste in Iceland and 1%-2% formic acid. The enzymatic breakdown and natural fermenting (natural bacterial flora in fish gut and

30

enzymes) fish silage is converted into a homogenous mince. To this about 10%-20% of algae, preferably Icelandic algae, may be added in geothermal water blend in the first 1-3 days after the initial fermentation and further allowed to ferment and enzymatically be converted into soluble nitrogen containing both the seaweed and soluble fish protein.

5

Compositions

Organic compositions of the present invention are water based and are formulated for a wide array of different dilutions with water in order to address a wide array of different applications. Dilution of compositions described herein is related to the particular

10 application. In general, a dosage of 1-25 ml organic composition/liter water is preferred as soil drench, seed treatment or foliar application; however, up to about fifty (50) ml organic composition of the present invention may be used per liter of water. Compositions of the present invention are capable of stimulating the proline-linked phenolic pathway in plants *and* acting as a bioenhancer at very low levels of 2-10 ml organic composition/liter of water.

15 Dilution of the organic compositions, for example, are in the range of 1-5 ml organic compositions/liter water. Higher concentrations of compositions of the present invention, e.g., in the range of 5-25 ml/liter water, are preferred for application to shrubs and trees.

Organic compositions of the present invention may be formulated to address different issues
20 faced by various growers. Natural extracts may be included in the compositions, for example, to facilitate and enhance to viability and growth of certain target species; however, other target species can be disadvantaged and thereby eliminated. Botanical extracts and complex polysaccharides Laminarin from Laminaria, for example, further increase the plant stimulating antioxidant properties of compositions of the invention. Other herbal extracts
25 from the family Lamiaceae used up to 5% w/v of organic compositions of the present invention, garlic extracts 5% extract of dry garlic extract in water (10 gram/100 ml extract) and use 1:20 dilution of the organic compositions of the present invention are added, for example, which, *inter alia*, increase anti-fungal properties. Neem extract, same as herbal extract, is further employed in compositions of the invention to increase natural anti pesticide
30 activity around the root zones. The activity can also be enhanced by other herbs added to it at 1% levels such as oregano, rosemary, thyme and other species in the botanical family Lamiaceae. For enhanced pesticide and antimicrobial properties of herbs, neem, onion waste and garlic extracts can be added at levels from 1%-5%.

Compositions described herein improve the growth and yield of agronomically important plants by the enhancement of root as well as shoot growth thus modifying the resulting structure of the plant. Organic compositions of the present invention are well suited for promoting the growth of flowers, fruits, vegetables, as well as shrubs and trees. The stimulation of the production of phenolic metabolites and aromatic compounds in treated plants leads to a significantly perceived improvement in the flavor and natural aroma of different crops. The phenotypic quality of treated flowering plants is also increased. Compositions of the present invention are also employed as a medium for seed germination (e.g., overnight soaking) and seed vigor stimulator. Compositions of the present invention may also be used, for example, as compost stimulator.

Bio-remediation

Compositions of the present invention moreover enrich treated soil directly and/or indirectly with beneficial microorganisms. This aspect of the present invention, for example, provides a means for significant improvement of soil symbiotic relationships including nitrogen fixing bacteria thus effecting soil improvement, particularly of marginal and dead soils.

Compositions described herein are particularly suitable for bio-remediation of flora habitation or regeneration otherwise of contaminated marginal or dead land (e.g., superfund sites) including but not limited to, for example, sites of petroleum, PCB, polyaromatic hydrocarbons, and/or dioxin contamination. Compositions of the present invention may be used, for example, to re-vegetate a contaminated area with a vegetation source such as Alfalfa and/or grasses and/or other monocot species. However, dicots, such as corn and other agronomic crops, may similarly be employed with compositions of the present invention, for example, to regenerate contaminated sites. As a corollary, the regeneration of plants in this manner, which demonstrate improved anti-oxidant response and hence detoxification of compounds, particularly carcinogenic agents, has significant impact on neutralizing and reducing the level and proximity of toxins.

Improved root growth

Compositions of the present invention promote the growth of beneficial soil bacteria in the plant root environment that in turn promote root development. Application of about 25-50 ml per liter of organic composition, for example, as a soil drench significantly enhanced the

population of soil microflora that promote plant growth. Treatment with compositions of the present invention also facilitates the association of beneficial bacteria in the root environment that inhibit the growth of plant pathogens. Treatment of plant soil with compositions of the present invention particularly facilitates and promotes the growth of microorganisms that
5 degrade organic matter and makes nutrients more easily available to plants. Organic compositions of the present invention also are employed to confer an improved regeneration of cuttings and new plants.

Organic compositions of the present invention may be used to benefit the production of fruits
10 including, for example, the transplant of strawberries and cranberry plants in bogs for enhanced growth and improved fruit quality. A dose of 2 ml of an organic composition of the present invention diluted in 1 liter of irrigation water, for example, has been demonstrated to enhance significantly the growth of strawberry and raspberry vegetative plants from tissue culture. Particularly, treatment at periods of watering during propagation of transplants
15 significantly enhances adaptation and survival. Strawberry varieties, for example, that normally exhibit poor transplanting characteristics have demonstrated an improvement of survival from 20% to 80% with treatment using the organic composition described herein. Increased widths of stems and roots, for example, are exhibited by transplanted plants treated with compositions of the present invention. Field adjustment of treated plants is moreover
20 improved using compositions of the present invention. Berry plants in the field, for example, may be treated using 10 ml of undiluted composition/liter of irrigation water per liter (e.g., sprayed on leaves every 30 days for strawberry). Berry plants, for example, may also be treated by means of irrigation. Raspberry and blackberry, for example, may be treated by means of administering about 25 ml per liter water to the soil every 30 days at an application
25 rate of about 400 to about 1000 liter per hectare. This technology is now suitable for all transplant seedlings going from greenhouse to field such as tomato and other transplant seedlings particularly where 99%-100% survival rate is required.

Improvement of vegetative and fruit yields

30 Soil application of about 50 ml organic composition of the present invention/liter of irrigation water at 30 day intervals with regular watering enhances the yield and size, *inter alia*, of tomato and egg plant, for example, between 15% to 20%, compared to absence of such application. The quality of the fruit and vegetables is generally better, i.e., fresher, with better

size, pigmentation, and taste. These benefits are expected with all types of vegetables, including cucumbers, squash, and melons, where preliminary results are promising.

5 An application of about 25 ml organic composition per liter of irrigation water is sprayed, for example, in the spring during early vegetative phase of perennial fruit stands such as cranberry. An application of about 10 ml per liter may be applied, for example, during the flowering phase and/or about 10 days prior to harvest. These applications improve pigmentation as well as fruit quality and yield.

10 An application of about 50 ml organic composition/liter of irrigation water may be employed to treat fruit orchards, for example, as a soil drench every 30 days during the growing season. Each application may be up to water holding capacity or an application rate of about 1000-1500 liter per hectare depending on soil type and temperature. In temperate regions for apples and temperate fruits an application rate can be about 1000 liter per hectare. Treatment
15 using about 1500 liters per hectare is suitable, for example, for citrus and Tropical and Sub-Tropical fruits.

Improvement of growth and stress-adaptation of transplant seedlings when moved from the greenhouse to an outdoor soil environment.

20 Leaf spray and/or soil-based drench application at a dose of about 2-5 ml organic composition/liter of irrigation water improves the survival of all greenhouse transplanted plants. Particularly, spray or soil drench and/or foliar application prior to transplanting enhances the strength of treated plants' stems through apparent increased lignification. Survival of transplants is significantly improved as well as conferred increased foliage and
25 aesthetic traits. In case of seeded transplants, treatment prior to germination in addition is very useful, of course, before application as a soil drench or foliar spray.

Improvement of seed vigor in seeded crops

Organic compositions of the present invention stimulate amino acid metabolism including
30 proline and stimulate plant growth regulators for improved vigor and robustness of the germinating seeds. Accordingly, organic compositions presented herein are successfully employed to enhance seed vigor during seed germination. Seeds are treated, for example, in a solution of about 2-5 ml organic composition per liter of irrigation water for 2-3 hours and

air-dried before germination. Appropriate volume may be readily determined based on the species-recommended seed rate and area of planting. Seed treatment generally translates into 10-20% improvement in germination rate and vigor than without the treatment. Pre-germination treatment is successful for all seeded plants, including, but not limited to cereals, legumes, vegetables and ornamentals. This is particularly preferred for germination of annuals, but can be employed to benefit the germination of large seeds as well. The use of compositions described herein is suitable for home gardens, particularly for growing vegetables.

10 **Enhancement of turf grass vegetative phase under cold conditions**

The organic compositions of the present invention are particularly useful to control water stress conditions and cold stress conditions. The soluble amino acids and peptides within compositions of the present invention stimulate light response and growth in plants. Treated vegetation is able to better grow, for example, in available light conditions. Application of about 10-25 ml organic composition per liter of irrigation water (higher dose in colder latitudes), for example, after mowing improves the vigor of lawn grass. Treatment stimulates rejuvenation of grass under cold spring conditions with much success. An application rate of about 400 liters per acre is recommended.

20 **Improvement of vegetative development and flowering in ornamentals**

Application of organic compositions of the invention as a foliar spray, for example, is particularly advantageous in promoting the vegetative development and flowering of ornamentals. Green vegetative phase with more branching and enhanced flowering can be expected when about 10-20 ml organic composition/liter of irrigation water is applied every 30 days, for example, during the growing season in gardens and outdoor field environments.

EXAMPLES

EXAMPLE I

Seed Treatment (for soft coated fruit and vegetable and other seeds)

- 30 a) Add 5 ml organic composition per one (1) liter of irrigation water.
- b) Soak seeds for 2-3 hours for soft coated seeds.
- c) Dry seeds in air in room temperature for 1-2 hours.

d) Sow the seeds as per seed producer's instructions.

e) Use the residual soak water as a soil drench and sprinkle over the soil after sowing.

For hard coated seeds, soak for 12 hours overnight in step b) and dry seeds in step c) for 2-5 hours before sowing.

5

EXAMPLE II

Improvement of Early Vegetative Growth (Green Phase 1)

- 10 a) Following germination use a dose of 5 ml organic composition per one (1) liter of irrigation water.
- b) Spray a rate of about 1000 liters per hectare or 400 liters per acre with a complete leaf drench. Adjust dosage for lesser area.
- c) Spray every 10 days until day 30.

For grasses, use above dosage after lawn mowing

15

EXAMPLE III

Improvement of Mid-Vegetative Phase (Green Phase 2)

- 20 a) Use a dose of 10 ml organic composition per liter of irrigation water.
- b) Spray a rate of about 1000 liters per hectare or 400 liters per acre with a complete leaf drench. Adjust dosage for lesser area.
- c) Spray every 10 days until day 60.
- d) Use dose of 25 ml per liter for shrubs, perennials and more woody stemmed species.

25

EXAMPLE IV

Improvement of Late-Vegetative Phase (Flowering and/or Seed Phase)

- 30 a) Use a dose of 10 ml organic composition per liter of irrigation water.
- b) Spray a rate of about 1000 liters per hectare or 400 liters per acre with a complete leaf drench. Adjust dosage for lesser area.
- c) Spray every 10 days until day 100.
- d) Use dose of 50 ml per liter for shrubs, perennials and more woody stemmed species.

EXAMPLE V**Trees and Perennial Potted Plants**

- a) Use dose of 50 ml organic composition per one (1) liter.
- b) Give one good soil soak up to water holding capacity (not excess) every 30 days from
5 May until September for out-door plants only.
- c) The remaining time when the soil is dry provide only regular water as needed.
- d) The general requirement is about 1 liter for a large ornamental tree with 20 inch pot
size and 100 ml to 250 ml for pot size of greater than 10 inches.
- e) Reduce dose to 25 ml per one (1) liter for pots less than 10 inches and reduce
10 application volume until is the soil is wet only up to water holding capacity and *not*
excess water.

Non-marine based Organic proteins from soy seed wastes and dairy whey proteins with final
nitrogen levels of 1% is available for Indoor Plants, where the application can be done every
45 days with regular watering at a dose of 25 ml per one (1) liter of irrigation water.

15

* * *

All publications and patents mentioned in the above specification are herein
incorporated by reference. Various modifications and variations of the described
20 compositions and methods of the invention will be apparent to those skilled in the art without
departing from the scope and spirit of the invention. Although the invention has been
described in connection with specific preferred embodiments, it should be understood that the
invention as claimed should not be unduly limited to such specific embodiments. Indeed,
various modifications of the described compositions and modes for carrying out the invention
25 which are obvious to those skilled in the art or related fields are intended to be within the
scope of the following claims.